

How partition function zeros help find out the finite-size scaling above the upper critical dimension

Tuesday, 16 January 2024 11:30 (20 minutes)

For many years the question of finite-size scaling above the upper critical dimension d_c was a source of many new theoretical results as well as new and interesting works in computer simulations, see [1] for a review. While in the thermodynamic limit system's scaling behaviour is mean-field, for the finite-size systems two scaling regimes are being considered. Simple approach is a trivial renormalization group fixed point, obtained by setting all fields to zero, called Gaussian fixed point, or G-scaling. Second scaling regime introduces a new scaling exponent $q = \max(1, d/d_c)$, that governs the scaling of correlation length ξ with system size L , it is called Q-scaling. The latter was confirmed for periodic boundary conditions, both at the infinite system's critical temperature and at the pseudocritical points. For lattices with free boundary conditions the shifting and rounding of the susceptibility peak that occurs at the pseudocritical point are big enough to position the pseudocriticality far from the critical point of the infinite system. This leads to both scaling regimes emerge, so theory predicted G-scaling at T_c and Q-scaling at T_L . Numerical validation of the finite-size scaling for free boundary conditions was not clear, as the boundary effects are too strong to observe the scaling picture accurately. In this work we show the shortcomings of the regular computer simulations methods and improve the quality of the finite-size scaling using the Lee-Yang zeros technique. Using the partition function zeros provides us with the more accurate scaling picture with smaller lattice sizes at hand [2].

1. Berche, B.; Ellis, T.; Holovatch, Yu.; Kenna, R. Phase Transitions above the Upper Critical Dimension. *SciPost Phys. Lect. Notes* 2022, 60. DOI: 10.21468/SciPostPhysLectNotes.60
2. Honchar Yu., Berche B., Holovatch Yu., Kenna R. When correlations exceed system size: finite-size scaling in free boundary conditions above the upper critical dimension. // preprint ArXiv: - 2023. - arXiv:2311.11721. (to appear in *Condens.Matter Phys* 2024, 1) DOI:10.48550/arXiv.2311.11721

Primary author: HONCHAR, Yulian (Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine, Lviv; Centre for Fluids and Complex Systems, Coventry University, Coventry CV1 5FB, UK; L4 Collaboration and Doctoral College for the Statistical Physics of Complex Systems, Lviv-Leipzig-Lorraine-Coventry, Europe)

Co-authors: BERCHE, Bertrand (Laboratoire de Physique et Chimie Théoriques, Université de Lorraine - CNRS, Nancy Cedex, France; L4 Collaboration and Doctoral College for the Statistical Physics of Complex Systems, Lviv-Leipzig-Lorraine-Coventry, Europe); HOLOVATCH, Yuriy (Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine, Lviv, 79000, Ukraine; Centre for Fluids and Complex Systems, Coventry University, Coventry CV1 5FB, UK; L4 Collaboration and Doctoral College for the Statistical Physics of Complex Systems, Lviv-Leipzig-Lorraine-Coventry, Europe; Complexity Science Hub Vienna, 1080 Vienna, Austria); KENNA, Ralph (Centre for Fluids and Complex Systems, Coventry University, Coventry CV1 5FB, UK; L4 Collaboration and Doctoral College for the Statistical Physics of Complex Systems, Lviv-Leipzig-Lorraine-Coventry, Europe)

Presenter: HONCHAR, Yulian (Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine, Lviv; Centre for Fluids and Complex Systems, Coventry University, Coventry CV1 5FB, UK; L4 Collaboration and Doctoral College for the Statistical Physics of Complex Systems, Lviv-Leipzig-Lorraine-Coventry, Europe)

Session Classification: Morning session 1

Track Classification: Condensed Matter Physics