Contribution ID: 101 Type: Oral

## The Painlevé I equation and Virasoro algebra

In this talk, I will explain how rank 5/2 representations of Virasoro algebra can be used to compute asymptotic expansion of the tau function of Painleve 1 equation near irregular singularity. The talk is based on recent paper of Hasmik Poghosyan and Rubik Poghossian, where they introduced conformal block with irregular vertex of rank 5/2 and conjectured that it is related to the partition function of  $H_0$  Argyres–Douglas theory. Additionally, I will present some improvements of this construction, which is the part of ongoing paper in collaboration with Oleg Lisovyy, Nikolai Iorgov and Kohei Iwaki.

**Primary author:** ZHURAVLOV, Yurii (Bogolyubov Institute for Theoretical Physics of the National Academy of Sciences of Ukraine)

**Co-authors:** IORGOV, Nikolai (Bogolyubov Institute for Theoretical Physics); LISOVYY, Oleg (Institut Denis-Poisson, Université de Tours); IWAKI, Kohei (The University of Tokyo)

**Presenter:** ZHURAVLOV, Yurii (Bogolyubov Institute for Theoretical Physics of the National Academy of Sciences of Ukraine)

Session Classification: MATHEMATICS

Track Classification: MATHEMATICS