Contribution ID: 63

Type: Oral

Stimulating exitable membrane of ORN with stochastic Markov process

Thursday, 26 September 2024 11:20 (20 minutes)

Excitable membrane of olfactory receptor neuron (ORN) is populated with up to several millions of identical receptor proteins (R) able to bind / release odor (O) molecules. The affinity of R to O depends on the odor presented, and this is the initial mechanism which is recuired for the olfactory selectivity to exist.

Figure 1: Simplified R structure. Modified from [3].

The affinity of R to O depends on the odor presented, and this is the very first step the olfactory selectivity builds up. Recently, [1], it was shown for a so called membrane-less ORN model that its selectivity can be much better than that of its R. A more realistic ORN model should include an excitable membrane with its electric transients. This introduces time parameter into the ORN's response to stimulation. The latter renders inappropriate used in [1] reasoning in terms of binomial distribution, and necessitates consideration of temporal properties of O binding-releasing and the membrane charging-discharging-firing.

In this contribution, we develop an approach in which the number of R bound with O n(t), is modeled as a Markov stochastic process. With each bound R,

as it is observed for insects [2], Fig. 1, we associate an open channel having conductance 0.015 nS, which injects a depolarizing current through the membrane.

The futher membrane evolution is governed by the leaky integrate-and-fire neuronal model, see Eq. (1).

$$c_M \frac{dV(t)}{dt} = -g_l(V(t) - V_{rest}) - n(t)g_R(V(t) - V_e),$$

where V(t) – is the membrane voltage; V_{rest} – is the resting voltage;

 c_M — is the total capacity of ORN's membrane;

 g_l — is the total leakage through it;

 $V_e-{\rm is}$ the reversal potential for current through open R;

 $\boldsymbol{n}(t)-\text{is the fluctuating number of open channels at moment <math display="inline">t$ due

to odor molecules bound with Rs;

 $g_R-{\rm is}$ the conductance of a single open channel.

A fast, very efficient method is developed for generating stochastic trajectories n(t) and solving Eq. (1) numerically, see Fig. 2. The first, introductory simulations based on this method, [4], support the conclusion made in [1] and before, that ORN's selectivity can be much better than that of its receptors R, provided that odors are presented in low concentrations.

Figure 2: An example of realization of stochastic process n(t) and corresponding membrane voltage, with three spikes emitted. Here, the total number of R per the ORN is $N = 2.5 \cdot 10^6$

[1] A. K. Vidybida, "Maximization of the olfactory receptor neuron selectivity in the sub-threshold regime", Ukrainian Journal of Physics, vol. 68, no.4, p. 266, 2023. Available: {\tt https://doi.org/10.15407/ujpe68.4.266}

 [2] K. Sato, M. Pellegrino, T. Nakagawa, T. Nakagawa, L. B. Vosshall, and
K. Touhara, "Insect olfactory receptors are heteromeric ligand-gated ion channels", Nature, vol. 452, no. 7190, pp. 1002–1006, 2008. Available: {\tt https://doi.org/10.1038/nature06850}

[3] D. Wicher, F. Miazzi, "Functional properties of insect olfactory receptors: ionotropic receptors and odorant receptors", Cell and Tissue Research, vol. 383(1):7-19 (2021). Available: {\tt https://doi.org/10.1007/s00441-020-03363-x}

[4] A. Vidybida, "Selectivity Gain in Olfactory Receptor Neuron at Optimal Odor Concentration," 2024 IEEE International Symposium on Olfaction and Electronic Nose (ISOEN), Grapevine, TX, USA, 2024, pp. 1-3. Available: {\tt https://ieeexplore.ieee.org/document/10556323}

Primary author: VIDYBIDA, Oleksandr (Bogolyubov Institute for Theoretical Physics)Presenter: VIDYBIDA, Oleksandr (Bogolyubov Institute for Theoretical Physics)Session Classification: Morning Session 3

Track Classification: STATISTICAL PHYSICS AND KINETIC THEORY