



QUANTUM CHEMICAL STUDY OF METHANOL INTERACTION WITH A PYRENE-BASED ACID

Kremen O. S.¹, Bychko I. B.², Nikolaienko T. Y.³, Lobanov V. V.¹, Strizhak P. E.²


¹*Chuiko Institute of Surface Chemistry of the NAS of Ukraine, Oleh Mudrak Str., 17, Kyiv-03164, Ukraine,*

²*L. V. Pisarzhevsky Institute of Physical Chemistry of the NAS of Ukraine, Prospect Nauki, 31, Kyiv-03028, Ukraine*


³*Taras Shevchenko National University of Kyiv, Faculty of Physics, Volodymyrska Str., 64/13, Kyiv 01601, Ukraine*

E-mail: kremenoksana@ukr.net

The conversion of methanol (MeOH) to dimethyl ether (DME) is a large-scale industrial process, and developing advanced catalysts to reduce costs is important. Oxygen-containing carbons are promising metal-free catalysts for many reactions. Therefore, this report presents quantum-chemical calculations (DFT, M062X/6-31G**/D3, ORCA 6.0.0) of the interaction between MeOH and pyrene-based acid (Pyr) in the gas phase at $T = 0$ K to elucidate the initial interaction mechanism for further investigation of the Methanol-to-DME conversion.



Pyr-COOH (A) $E_{\text{total}} -804.02749406$ Eh



Pyr-COOH (B) $E_{\text{total}} -804.02976006$ Eh

The distance between atoms in Å, the O-C-C dihedral angles in degrees (denoted in green), the atomic charge ($|e|$) by Mulliken in bold. S(6) and S(5) – motifs in the graph-set assignments according to [1]. ● – H, ● – C, ● – O.


Geometry optimisation yielded two acid isomers (A and B), depending on the orientation of the –COOH group plane relative to the aromatic plane. The total energy in a.u. The E_{total} of the planar B-Pyr acid is lower by 5.95 kJ/mol than that of the twisted A-Pyr acid molecule.

The topological analysis of the electron density distribution (Bader's Quantum Theory of Atoms in Molecules) confirmed the presence of two intramolecular hydrogen bonds (HBs) in the B-Pyr as evidenced by the presence of the corresponding bond critical points (BCPs) of type (3, –1). In contrast, only a single intramolecular HB was observed in the A-Pyr acid. Despite the $\text{C}=\text{O} \cdots \text{H}-\text{C}$ length being 2.361 Å (less than the sum of the van der Waals radii of the O and H atoms, which is 2.5 Å), there are no BCPs according to Bader's analysis.


Two possible ways of MeOH interaction with the B-Pyr acid are considered.

(1) Interaction of Methanol with a Pyrene-Based Acid: Formation of a Cyclic Complex (CC)


The CC is stabilised by two intermolecular HBs, forming a six-membered ring (6MR) between the –COOH group atoms of the Pyr acid and the –OH group atoms of the MeOH molecule. The interaction energy is –66.48 kJ/mol. The CC have two tautomeric forms (causing asymmetry in the carbon skeleton), more twisted CCa , and CCb , which can interconvert through an activation barrier of 42.40 kJ/mol. The transition state (TS) for the transition between CCa , and CCb structural isomers is characterised by a flip-flop switching of a hydrogen atom between adjacent acceptor centres (two O atoms of the –COOH group and one O atom of the MeOH), forming a two-sided hydrogen trap. The configuration of CCb is thermodynamically more stable than CCa , since its formation energy is lower by 4.43 kJ/mol.



Cyclic Complex a (CCa)



Transition state (TS)




Cyclic Complex b (CCb)

(2) Interaction of Methanol with a Pyrene-Based Acid: Attack of the C=O of the Pyr by MeOH

Attack on the $\text{C}=\text{O}$ bond of the Pyr leads to the formation of a tetrahedral structure because the carbon atom orbitals in the –COOH group change hybridisation from sp^2 to sp^3 . This atom is bound to four substituents: a Pyr-based group ($-\text{C}_{16}\text{H}_9$), two –OH groups and a single methoxy group ($-\text{OCH}_3$). The analysis of the atomic vibrations for the cyclic four-membered TS structure confirmed the presence of a single imaginary frequency at $i1620.70$ cm^{-1} . The activation barrier is +169.01 kJ/mol, and the formation energy is +23.88 kJ/mol.

Based on the obtained energetic parameters, it was established that the bonding of the MeOH molecule preferentially proceeds via the formation of 6MR CC structures, as this pathway is significantly more energetically favourable than direct attack on the $\text{C}=\text{O}$ bond of the –COOH group.



Tetrahedral structure

This research was partially funded by the National Research Foundation of Ukraine (grant 2023.03/0041).

[1]. M. C. Etter, Encoding and decoding hydrogen-bond patterns of organic compounds, Acc. Chem. Res., 23 (1990) 120–126, <https://doi.org/10.1021/ar00172a005>.